

# Hi-Moisture Corn Mixing Chart



**Directions:** Add some water to the applicator tank, then add the required amount of inoculant and continue filling the tank.

- **One Tablespoon of inoculant is 10 grams.**
- **One grams of inoculant treats 1 ton/34 bushels of HM corn.**
- **One Tablespoon, with 5 gallons of water treats 10 tons/340 bushels of high moisture corn with an application rate of 1 gallon to 68 bushels.**
- When in doubt, always add more inoculant. Extra coverage will not damage the crop.
- Please, see charts below.
- These directions do not apply to Pro-Siler or Dohrmann Applicators. See other PDF for Pro-Siler/Dohrmann mixing.

25 Gallon Tank		
Amount of Inoculant	Gallons of Water	Application Rate
50 ton pack	25	1 gal. to 68 bu.
100 ton pack	25	1 gal. to 135 bu.

55 Gallon Tank		
Amount of Inoculant	Gallons of Water	Application Rate
50 ton pack	55	1 gal. to 30 bu.
100 ton pack	55	1 gal. to 60 bu.

## More Info

- Diluting inoculant in a larger volume of water ensures maximum coverage.
- Do not use chlorinated water.
- Do not use hot water.
- For best results use in a sprayer that has a recirculating pump.
- Use only FLOOD jets.
- Use NO SCREENS.
- Use any leftover mixture within 48 hours after mixing.
- Run clear water through applicator after use.
- Store package in a cool dry area.

**Any Questions?**

**Call: 920-251-5916**

[RK@strongmicrobials.com](mailto:RK@strongmicrobials.com)

**StrongMicrobials.com**

Made in Milwaukee, WI