

Corn Silage Mixing Chart



Directions: Add some water to the applicator tank, then add the required amount of inoculant and continue filling the tank.

- **One Tablespoon of inoculant is 10 grams.**
- **0.5 grams of inoculant treats 1 ton of corn silage.**
- **One Tablespoon of inoculant treats 20 tons of corn silage.**
- When in doubt, always add more inoculant. Extra coverage will not damage the crop.
- Please, see charts below for wagon size.
- These directions do not apply to Pro-Siler or Dohrmann Applicators. See other PDF for Pro-Siler/Dohrmann mixing

6 Ton Wagon Loads	
Number of Loads	Tablespoons of Inoculant
5	1½
10	3
15	4½
20	6
25	7½

8 Ton Wagon Loads	
Number of Loads	Tablespoons of Inoculant
5	2
10	4
15	6
20	8
25	10

10 Ton Wagon Loads	
Number of Loads	Tablespoons of Inoculant
5	2½
10	5
15	7½
20	10
25	12½

12 Ton Wagon Loads	
Number of Loads	Tablespoons of Inoculant
5	3
10	6
15	9
20	12
25	15

More Info

- Diluting inoculant in a larger volume of water ensures maximum coverage.
- Do not use chlorinated water.
- Do not use hot water.
- For best results use in a sprayer that has a recirculating pump.
- Use only FLOOD jets.
- Use NO SCREENS.
- Use any leftover mixture within 48 hours after mixing.
- Run clear water through applicator after use.
- Store package in a cool dry area.

Any Questions?

Call: 920-251-5916

RK@strongmicrobials.com

StrongMicrobials.com

Made in Milwaukee, WI